AUTHOR INDEX VOLUME 132

A

Abe, H., Obinata, T., Minamide, L.S., and Bamburg, J.R. *Xenopus laevis* actin-depolymerizing factor/cofilin: A phosphorylation-regulated protein essential for development. 871.

Aebi, U. *See* Goulielmos, Gounari, Remington, Müller, Häner, Aebi, and Georgatos.

Aizawa, H., Sutoh, K., and Yahara, I. Overexpression of cofilin stimulates bundling of actin filaments, membrane ruffling, and cell movement in *Dictyostelium*. 335.

Aizawa, S. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.

Alexander, C.M. *See* Sympson, Talhouk, Alexander, Chin, Clift, Bissell, and Werb.

Allen, T.StC. *See* Myers, Goh, Allen, Bucher, and Bogaert.

Altruda, F. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.

Andrés, V., and Walsh, K. Myogenin expression, cell cycle withdrawal, and phenotypic differentiation are temporally separable events that precede cell fusion upon myogenesis. 657.

Arai, K. *See* Yoshida, Fujisawa-Sehara, Taki, Arai, and Nabeshima.

Aruffo, A. *See* Bartolazzi, Nocks, Aruffo, Spring, and Stamenkovic.

B

Bacarra, A.E. *See* Parkos, Colgan, Liang, Nusrat, Bacarra, Carnes, and Madara.

Bachant, J.B., and Klymkowsky, M.W. A nontetrameric species is the major soluble form of keratin in *Xenopus* oocytes and rabbit reticulocyte lysates. 153.

Baeuerle, P.A. *See* Pahl, Sester, Burgert, and Baeuerle.

Bajer, A.S. *See* Khodjakov, Cole, Bajer, and Rieder.

Balzac, F. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.

Bamburg, J.R. *See* Abe, Obinata, Minamide, and Bamburg.

Bartolazzi, A., Nocks, A., Aruffo, A., Spring, F., and Stamenkovic, I. Glycosylation of CD44 is implicated in CD44-mediated cell adhesion to hyaluronan. 1199.

Becker, E. *See* Cabrera, Díaz-Rodríguez, Becker, Martín-Zanca, and Pandiella.

Becker, J.C. *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.

Belkin, A.M., Zhidkova, N.I., Balzac, F., Altruda, F., Tomatis, D., Maier, A., Tarone, G., Koteliansky, V.E., and Burridge, K. β1D integrin displaces the β1A isoform in striated muscles: Localization at junctional structures and signaling potential in nonmuscle cells. 211.

Benedetti, E.L. *See* Dunia, Smit, van der Valk, Bloemendal, Borst, and Benedetti.

Benian, G.M., Tinley, T.L., Tang, X., and Borodovsky, M. The *Caenorhabditis elegans* gene *unc-89*, required for muscle M-line assembly, encodes a giant modular protein composed of Ig and signal transduction domains. 835.

Beug, H. *See* Fialka, Schwarz, Reichmann, Oft, Busslinger, and Beug.

Bissell, M.J. *See* Sympson, Talhouk, Alexander, Chin, Clift, Bissell, and Werb.

Blobel, C.P. *See* Weskamp, Krätzschmar, Reid, and Blobel.

Block, G.D., Locker, J., Bowen, W.C., Petersen, B.E., Katyal, S., Strom, S.C., Riley, T., Howard, T.A., and Michalopoulos, G.K. Population expansion, clonal growth, and specific differentiation patterns in primary cultures of hepatocytes induced by HGF/SF, EGF and TGFα in a chemically defined (HGM) medium. 1133.

Bloemendal, H. *See* Dunia, Smit, van der Valk, Bloemendal, Borst, and Benedetti.

Bogaert, T. *See* Myers, Goh, Allen, Bucher, and Bogaert.

Borisy, G.G. *See* Wang, Peloquin, Zhai, Bulinski, and Borisy.

Borodovsky, M. *See* Benian, Tinley, Tang, and Borodovsky.

Borst, P. *See* Dunia, Smit, van der Valk, Bloemendal, Borst, and Benedetti.

Bos-Mikich, A. *See* Kono, Jones, Bos-Mikich, Whittingham, and Carroll.

Bowen, W.C. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.

Bravo, N.S. *See* Paladini, Takahashi, Bravo, and Coulombe.

Brodsky, J.L. *See* McCracken and Brodsky.

Bron, R. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.

Bucher, E.A. *See* Myers, Goh, Allen, Bucher, and Bogaert.

Bulinski, J.C. *See* Wang, Peloquin, Zhai, Bulinski, and Borisy.

Burgert, H.-G. *See* Pahl, Sester, Burgert, and Baeuerle.

Burgeson, R.E. *See* Champliaud, Lunstrum, Rousselle, Nishiyama, Keene, and Burgeson.

Burridge, K. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.

Busslinger, M. *See* Fialka, Schwarz, Reichmann, Oft, Busslinger, and Beug.

C

Cabrera, N., Díaz-Rodríguez, E., Becker, E., Martín-Zanca, D., and Pandiella, A. TrkA receptor ectodomain cleavage generates a tyrosine-phosphorylated cell-associated fragment. 427.

Caffarelli, A.D. *See* Oda, Scott, Hefner-Gravink, Caffarelli, and Klionsky.

Carnes, D.K. *See* Parkos, Colgan, Liang, Nusrat, Bacarra, Carnes, and Madara.

Carroll, J. *See* Kono, Jones, Bos-Mikich, Whittingham, and Carroll.

Carter, W.G. *See* Xia, Gil, and Carter.

Champliaud, M.-F., Lunstrum, G.P., Rousselle, P., Nishiyama, T., Keene, D.R., and Burgeson, R.E. Human amnion contains a novel laminin variant, laminin 7, which like laminin 6, covalently associates with laminin 5 to promote stable epithelial–stromal attachment. 1189.

Chan, P.-Y. *See* Dustin, Ferguson, Chan, Springer, and Golan.

Cheresh, D.A. *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.

Chin, J.R. *See* Sympson, Talhouk, Alexander, Chin, Clift, Bissell, and Werb.

Clark, R.A.F. *See* Xu and Clark.

Clase, K.L. *See* Hannon, Kudla, McAvoy, Clase, and Olwin.

Clayton, C. *See* Häusler, Stierhof, Wirtz, and Clayton.

Clement, A. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.

Clemson, C.M., McNeil, J.A., Willard, H.F., and Lawrence, J.B. XIST RNA paints the inactive X chromosome at interphase: Evidence for a novel RNA involved in nuclear/chromosome structure. 259.

Clift, S.M. *See* Sympson, Talhouk, Alexander, Chin, Clift, Bissell, and Werb.

Cole, D.G. *See* Wedaman, Meyer, Rashid, Cole, and Scholey.

Cole, R.W. *See* Khodjakov, Cole, Bajer, and Rieder.

Colgan, S.P. *See* Parkos, Colgan, Liang, Nusrat, Bacarra, Carnes, and Madara.

Condon, P.J. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.

Cooper, J.A. *See* Waddle, Karpova, Waterston, and Cooper.
Cooper, J.B. *See* Prekeris, Mayhew, Cooper, and Terrian.
Corvera, S. *See* Shpetner, Joly, Hartley, and Corvera.
Coulombe, P.A. *See* Paladini, Takahashi, Bravo, and Coulombe.
Cowan, N.J. *See* Lewis, Tian, Vainberg, and Cowan.
Creutz, C.E., Snyder, S.L., Daigle, S.N., and Redick, J. Identification, localization, and functional implications of an abundant nematode annexin. 1079.

D

Daigle, S.N. *See* Creutz, Snyder, Daigle, and Redick.
Daou, M.-C. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.
David, G. *See* Mertens, van der Schueren, van den Berghe, and David.
Davies, P. *See* Vincent, Rosado, and Davies.
Davis, T.N. *See* Friedman, Sundberg, Huang, and Davis.
Deiner, M. *See* Ferns, Deiner, and Hall.
Derynck, R. *See* Erlebacher and Derynck.
de Vree, W.J.A. *See* Koolwijk, van Erck, de Vree, Vermeer, Weich, Hanemaaijer, and van Hinsbergh.
Díaz-Rodríguez, E. *See* Cabrera, Díaz-Rodríguez, Becker, Martín-Zanca, and Pandiella.
Dice, J.F. *See* Hayes and Dice.
Dietrich, J., Neisig, A., Hou, X., Wegener, A.-M.K., Gajhede, M., and Geisler, C. Role of CD3γ in T cell receptor assembly. 299.
Dittié, A.S., Hajibagheri, N., and Tooze, S.A. The AP-1 adaptor complex binds to immature secretory granules from PC12 cells, and is regulated by ADP-ribosylation factor. 523.
Donaldson, A.D., and Kilmartin, J.V. Spc42p: A phosphorylated component of the *S. cerevisiae* spindle pole body (SPB) with an essential function during SPB duplication. 887.
Downey, G.P. *See* Fukushima, Waddell, Grinstein, Goss, Orlowski, and Downey.
Dunia, I., Smit, J.J.M., van der Valk, M.A., Bloemendal, H., Borst, P., and Benedetti, E.L. Human multidrug resistance 3-P-glycoprotein expression in transgenic mice induces lens membrane alterations leading to cataract. 701.
Dustin, M.L., Ferguson, L.M., Chan, P.-Y., Springer, T.A., and Golan, D.E. Visualization of CD2 interaction with LFA-3 and determination of the two-dimensional dissociation constant for adhesion receptors in a contact area. 465.

E

Eady, R.A.J. *See* Porter, Leitgeb, Melton, Swensson, Eady, and Magin.
Echeverri, C.J., Paschal, B.M., Vaughan, K.T., and Vallee, R.B. Molecular characterization of the 50-kD subunit of dynactin reveals function for the complex in chromosome alignment and spindle organization during mitosis. 617.
Egelhoff, T.T. *See* Kolman, Futey, and Egelhoff.
Ekblom, P. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Erlebacher, A., and Derynck, R. Increased expression of TGF-β2 in osteoblasts results in an osteoporosis-like phenotype. 195.
Ernst, J.D. *See* Zimmerli, Majeed, Gustavsson, Stendahl, Sanan, and Ernst.

F

Fagotto, F., Funayama, N., Glück, U., and Gumbiner, B.M. Binding to cadherins antagonizes the signaling activity of β-catenin during axis formation in *Xenopus*. 1105.
Faissner, A. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Fares, H., Goetsch, L., and Pringle, J.R. Identification of a developmentally regulated septin and involvement of the septins in spore formation in *Saccharomyces cerevisiae*. 399.
Fässler, R. *See* Wennerberg, Lohikangas, Gullberg, Pfaff, Johansson, and Fässler.
Ferguson, L.M. *See* Dustin, Ferguson, Chan, Springer, and Golan.
Ferns, M., Deiner, M., and Hall, Z. Agrin-induced acetylcholine receptor clustering in mammalian muscle requires tyrosine phosphorylation. 937.
Fialka, I., Schwarz, H., Reichmann, E., Oft, M., Busslinger, M., and Beug, H. The estrogen-dependent c-JunER protein causes a reversible loss of mammary epithelial cell polarity involving a destabilization of adherens junctions. 1115.
Forbes, D.J. *See* Macaulay and Forbes.
Frank, R. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Friedman, D.B., Sundberg, H.A., Huang, E.Y., and Davis, T.N. The 110-kD spindle pole body component of *Saccharomyces cerevisiae* is a phosphoprotein that is modified in a cell cycle–dependent manner. 903.
Fujisawa-Sehara, A. *See* Yoshida, Fujisawa-Sehara, Taki, Arai, and Nabeshima.
Fukushima, T., Waddell, T.K., Grinstein, S., Goss, G.G., Orlowski, J., and Downey, G.P. Na^+/H^+ exchange activity during phagocytosis in human neutrophils: Role of Fcγ receptors and tyrosine kinases. 1037.
Funakoshi, T. *See* Hirokawa, Funakoshi, Sato-Harada, and Kanai.
Funayama, N. *See* Fagotto, Funayama, Glück, and Gumbiner.
Futey, L.M. *See* Kolman, Futey, and Egelhoff.
Futter, C.E., Pearse, A., Hewlett, L.J., and Hopkins, C.R. Multivesicular endosomes containing internalized EGF–EGF receptor complexes mature and then fuse directly with lysosomes. 1011.

G

Gajhede, M. *See* Dietrich, Neisig, Hou, Wegener, Gajhede, and Geisler.
Geisler, C. *See* Dietrich, Neisig, Hou, Wegener, Gajhede, and Geisler.
Georgatos, S.D. *See* Goulielmos, Gounari, Remington, Müller, Häner, Aebi, and Georgatos.
Geuze, H.J. *See* Glickman, Morton, Slot, Kornfeld, and Geuze.
— *See* Stoorvogel, Oorschot, and Geuze.
Gil, S.G. *See* Xia, Gil, and Carter.
Glickman, J.N., Morton, P.A., Slot, J.W., Kornfeld, S., and Geuze, H.J. The biogenesis of the MHC class II compartment in human I-Cell disease B lymphoblasts. 769.
Glück, U. *See* Fagotto, Funayama, Glück, and Gumbiner.
Goetsch, L. *See* Fares, Goetsch, and Pringle.
Goh, P.-Y. *See* Myers, Goh, Allen, Bucher, and Bogaert.
Golan, D.E. *See* Dustin, Ferguson, Chan, Springer, and Golan.
Gorza, L. *See* Rossetto, Gorza, Schiavo, Schiavo, Scheller, and Montecucco.
Goss, G.G. *See* Fukushima, Waddell, Grinstein, Goss, Orlowski, and Downey.
Götz, B., Scholze, A., Clement, A., Joester, A., Schütte, K., Wigger, F., Frank, R., Spiess, E., Ekblom, P., and Faissner, A. Tenascin-C contains distinct adhesive, anti-adhesive, and neurite outgrowth promoting sites for neurons. 681.
Goulielmos, G., Gounari, F., Remington, S., Müller, S., Häner, M., Aebi, U., and Georgatos, S.D. Filensin and phakinin form a novel type of beaded intermediate filaments and coassemble de novo in cultured cells. 643.
Gounari, F. *See* Goulielmos, Gounari, Remington, Müller, Häner, Aebi, and Georgatos.
Grinstein, S. *See* Fukushima, Waddell, Grinstein, Goss, Orlowski, and Downey.
Grote, E., and Kelly, R.B. Endocytosis of VAMP is facilitated by a synaptic vesicle targeting signal. 537.
Gullberg, D. *See* Wennerberg, Lohikangas, Gullberg, Pfaff, Johansson, and Fässler.
Gumbiner, B.M. *See* Fagotto, Funayama, Glück, and Gumbiner.
Gustavsson, M. *See* Zimmerli, Majeed, Gustavsson, Stendahl, Sanan, and Ernst.

H

Haggarty, B.S. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Hajibagheri, N. *See* Dittié, Hajibagheri, and Tooze.
Hall, Z. *See* Ferns, Deiner, and Hall.
Hamada, H. *See* Okazawa, Shimizu, Kamei, Imafuku, Hamada, and Kanazawa.
Hamamoto, S. *See* Lupashin, Hamamoto, and Schekman.
Hanemaaijer, R. *See* Koolwijk, van Erck, de Vree, Vermeer, Weich, Hanemaaijer, and van Hinsbergh.
Häner, M. *See* Goulielmos, Gounari, Remington, Müller, Häner, Aebi, and Georgatos.
Hannon, K., Kudla, A.J., McAvoy, M.J., Clase, K.L., and Olwin, B.B. Differentially expressed fibroblast growth factors regulate skeletal muscle development through autocrine and paracrine mechanisms. 1151.
Hanzawa, N. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Harris, S.L., and Waters, M.G. Localization of a yeast early Golgi mannosyltransferase, Och1p, involves retrograde transport. 985.
Hart, T.K. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Hartley, D. *See* Shpetner, Joly, Hartley, and Corvera.
Häusler, T., Stierhof, Y.-D., Wirtz, E., and Clayton, C. Import of a DHFR hybrid protein into glycosomes in vivo is not inhibited by the folate-analogue aminopterin. 311.
Hayes, S.A., and Dice, J.F. Roles of molecular chaperones in protein degradation. 255.
Hefner-Gravink, A. *See* Oda, Scott, Hefner-Gravink, Caffarelli, and Klionsky.
Hewlett, L.J. *See* Futter, Pearse, Hewlett, and Hopkins.
Hirokawa, N., Funakoshi, T., Sato-Harada, R., and Kanai, Y. Selective stabilization of tau in axons and microtubule-associated protein 2C in cell bodies and dendrites contributes to polarized localization of cytoskeletal proteins in mature neurons. 667.
Hisanaga, S.-I. *See* Okumura, Sekiai, Hisanaga, Tachibana, and Kishimoto.
Hojo, M. *See* Rajasekaran, Hojo, Huima, and Rodriguez-Boulan.
Hopkins, C.R. *See* Futter, Pearse, Hewlett, and Hopkins.
Hou, X. *See* Dietrich, Neisig, Hou, Wegener, Gajhede, and Geisler.
Howard, T.A. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Hoxie, J.A. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Huang, E.Y. *See* Friedman, Sundberg, Huang, and Davis.
Huima, T. *See* Rajasekaran, Hojo, Huima, and Rodriguez-Boulan.
Hynes, J.B. *See* Rijnboutt, Jansen, Posthuma, Hynes, Schornagel, and Strous.

I

Imafuku, I. *See* Okazawa, Shimizu, Kamei, Imafuku, Hamada, and Kanazawa.
Inagaki, M. *See* Sekimata, Tsujimura, Tanaka, Takeuchi, Inagaki, and Inagaki.

J

Jansen, G. *See* Rijnboutt, Jansen, Posthuma, Hynes, Schornagel, and Strous.
Joester, A. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Johansson, S. *See* Wennerberg, Lohikangas, Gullberg, Pfaff, Johansson, and Fässler.
Joly, M. *See* Shpetner, Joly, Hartley, and Corvera.
Jones, K.T. *See* Kono, Jones, Bos-Mikich, Whittingham, and Carroll.

K

Kamei, M. *See* Okazawa, Shimizu, Kamei, Imafuku, Hamada, and Kanazawa.
Kanai, Y. *See* Hirokawa, Funakoshi, Sato-Harada, and Kanai.
Kanazawa, I. *See* Okazawa, Shimizu, Kamei, Imafuku, Hamada, and Kanazawa.
Kanwar, Y.S. Wada, Kumar, Liu, Ruoslahti, Reichardt, Marvaldi, and Kanwar.
Kaplan, D.R. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.
Karpova, T.S. *See* Waddle, Karpova, Waterston, and Cooper.
Katyal, S. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Keen, J.H. *See* Santini and Keen.
Keene, D.R. *See* Champliaud, Lunstrum, Rousselle, Nishiyama, Keene, and Burgeson.
Kelly, R.B. *See* Grote and Kelly.
Khodjakov, A., Cole, R.W., Bajer, A.S., and Rieder, C.L. The force for poleward chromosome motion in *Haemanthus* cells acts along the length of the chromosome during metaphase but only at the kinetochore during anaphase. 1093.
Kilmartin, J.V. *See* Donaldson and Kilmartin.
Kishimoto, T. *See* Okumura, Sekiai, Hisanaga, Tachibana, and Kishimoto.
Klionsky, D.J. *See* Oda, Scott, Hefner-Gravink, Caffarelli, and Klionsky.
Klymkowsky, M.W. *See* Bachant and Klymkowsky.
Kolman, M.F., Futey, L.M., and Egelhoff, T.T. *Dictyostelium* myosin heavy chain kinase A regulates myosin localization during growth and development. 101.
Kono, T., Jones, K.T., Bos-Mikich, A., Whittingham, D.G., and Carroll, J. A cell cycle–associated change in Ca^{2+} releasing activity leads to the generation of Ca^{2+} transients in mouse embryos during the first mitotic division. 915.
Koolwijk, P., van Erck, M.G.M., de Vree, W.J.A., Vermeer, M.A., Weich, H.A., Hanemaaijer, R., and van Hinsbergh, V.W.M. Cooperative effect of TNFα, bFGF, and VEGF on the formation of tubular structures of human microvascular endothelial cells in a fibrin matrix. Role of urokinase activity. 1177.
Kornfeld, S. *See* Glickman, Morton, Slot, Kornfeld, and Geuze.
— *See* Rohrer, Schweizer, Russell, and Kornfeld.
— *See* Schweizer, Kornfeld, and Rohrer.
Koteliansky, V.E. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.
Krätzschmar, J. *See* Weskamp, Krätzschmar, Reid, and Blobel.
Kudla, A.J. *See* Hannon, Kudla, McAvoy, Clase, and Olwin.
Kumar, A. *See* Wada, Kumar, Liu, Ruoslahti, Reichardt, Marvaldi, and Kanwar.

L

LaBranche, C.C. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Lachyankar, M.B. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.
Landmesser, L. *See* Rafuse and Landmesser.
Lawrence, J.B. *See* Clemson, McNeil, Willard, and Lawrence.
Lazarow, P.B. *See* Zhang and Lazarow.
Lebakken, C.S., and Rapraeger, A.C. Syndecan-1 mediates cell spreading in transfected human lymphoblastoid (Raji) cells. 1209.
Lee, W.M.F. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Lefebvre, P.A. *See* Smith and Lefebvre.
Leitgeb, S. *See* Porter, Leitgeb, Melton, Swensson, Eady, and Magin.
Lemmon, V.P. *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.
Lewis, S.A., Tian, G., Vainberg, I.E., and Cowan, N.J. Chaperonin-mediated folding of actin and tubulin. 1.
Li, L., Miano, J.M., Mercer, B., and Olson, E.N. Expression of the *SM22α* promoter in transgenic mice provides evidence for distinct transcriptional regulatory programs in vascular and visceral smooth muscle cells. 849.

Liang, T. *See* Parkos, Colgan, Liang, Nusrat, Bacarra, Carnes, and Madara.
Liu, Z. *See* Wada, Kumar, Liu, Ruoslahti, Reichardt, Marvaldi, and Kanwar.
Locker, J. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Lohikangas, L. *See* Wennerberg, Lohikangas, Gullberg, Pfaff, Johansson, and Fässler.
Lopes-Cardozo, M. *See* van der Bijl, Lopes-Cardozo, and van Meer.
Lunstrum, G.P. *See* Champliaud, Lunstrum, Rousselle, Nishiyama, Keene, and Burgeson.
Lupashin, V.V., Hamamoto, S., and Schekman, R.W. Biochemical requirements for the targeting and fusion of ER-derived transport vesicles with purified yeast Golgi membranes. 277.
— *See* Sapperstein, Lupashin, Schmitt, and Waters.
Lyon, M.K. *See* Marr, Mastronarde, and Lyon.

M

Macaulay, C., and Forbes, D.J. Assembly of the nuclear pore: Biochemically distinct steps revealed with NEM, GTPγS, and BAPTA. 5.
Madara, J.L. *See* Parkos, Colgan, Liang, Nusrat, Bacarra, Carnes, and Madara.
Magin, T.M. *See* Porter, Leitgeb, Melton, Swensson, Eady, and Magin.
Maier, A. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.
Majeed, M. *See* Zimmerli, Majeed, Gustavsson, Stendahl, Sanan, and Ernst.
Marr, K.M., Mastronarde, D.N., and Lyon, M.K. Two-dimensional crystals of photosystem II: Biochemical characterization, cryo-electron microscopy and localization of the D1 and cytochrome b559 polypeptides. 823.
Marsh, M. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Martín-Zanca, D. *See* Cabrera, Díaz-Rodríguez, Becker, Martín-Zanca, and Pandiella.
Marvaldi, J. *See* Wada, Kumar, Liu, Ruoslahti, Reichardt, Marvaldi, and Kanwar.
Mastronarde, D.N. *See* Marr, Mastronarde, and Lyon.
Mayhew, M.W. *See* Prekeris, Mayhew, Cooper, and Terrian.
McAvoy, M.J. *See* Hannon, Kudla, McAvoy, Clase, and Olwin.
McCracken, A.A., and Brodsky, J.L. Assembly of ER-associated protein degradation in vitro: Dependence on cytosol, calnexin, and ATP. 291.
McKinnon, C.A. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.
McNeil, J.A. *See* Clemson, McNeil, Willard, and Lawrence.
Melton, D.W. *See* Porter, Leitgeb, Melton, Swensson, Eady, and Magin.
Mercer, B. *See* Li, Miano, Mercer, and Olson.
Mertens, G., van der Schueren, B., van den Berghe, H., and David, G. Heparan sulfate expression in polarized epithelial cells: The apical sorting of glypican (GPI-anchored proteoglycan) is inversely related to its heparan sulfate content. 487.
Meyer, D.W. *See* Wedaman, Meyer, Rashid, Cole, and Scholey.
Miano, J.M. *See* Li, Miano, Mercer, and Olson.
Michalopoulos, G.K. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Minamide, L.S. *See* Abe, Obinata, Minamide, and Bamburg.
Miyauchi, Y. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Mondésert, G., and Reed, S.I. *BED1*, a gene encoding a galactosyltransferase homologue, is required for polarized growth and efficient bud emergence in *Saccharomyces cerevisiae*. 137.
Montecucco, C. *See* Rossetto, Gorza, Schiavo, Schiavo, Scheller, and Montecucco.
Montgomery, A.M.P., Becker, J.C., Siu, C.-H., Lemmon, V.P., Cheresh, D.A., Pancook, J.D., Zhao, X., and Reisfeld, R.A. Human neural cell adhesion molecule L1 and rat homologue NILE are ligands for integrin $\alpha_v\beta_3$. 475.
Morton, P.A. *See* Glickman, Morton, Slot, Kornfeld, and Geuze.
Müller, S. *See* Goulielmos, Gounari, Remington, Müller, Häner, Aebi, and Georgatos.
Myers, C.D., Goh, P.-Y., Allen, T.StC., Bucher, E.A., and Bogaert, T. Developmental genetic analysis of troponin T mutations in striated and nonstriated muscle cells of *Caenorhabditis elegans*. 1061.

N

Nabeshima, Y. *See* Yoshida, Fujisawa-Sehara, Taki, Arai, and Nabeshima.
Neisig, A. *See* Dietrich, Neisig, Hou, Wegener, Gajhede, and Geisler.
Ng, D.T.W., and Walter, P. ER membrane protein complex required for nuclear fusion. 499.
Nishikawa, S.-I. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Nishiyama, T. *See* Champliaud, Lunstrum, Rousselle, Nishiyama, Keene, and Burgeson.
Nocks, A. *See* Bartolazzi, Nocks, Aruffo, Spring, and Stamenkovic.
Nusrat, A. *See* Parkos, Colgan, Liang, Nusrat, Bacarra, Carnes, and Madara.

O

Obinata, T. *See* Abe, Obinata, Minamide, and Bamburg.
Oda, M.N., Scott, S.V., Hefner-Gravink, A., Caffarelli, A.D., and Klionsky, D.J. Identification of a cytoplasm to vacuole targeting determinant in aminopeptidase I. 999.
Oft, M. *See* Fialka, Schwarz, Reichmann, Oft, Busslinger, and Beug.
Oh, Y.-K., and Swanson, J.A. Different fates of phagocytosed particles after delivery into macrophage lysosomes. 585.
Okazawa, H., Shimizu, J., Kamei, M., Imafuku, I., Hamada, H., and Kanazawa, I. Bcl-2 inhibits retinoic acid–induced apoptosis during the neural differentiation of embryonal stem cells. 955.
Okumura, E., Sekiai, T., Hisanaga, S.-I., Tachibana, K., and Kishimoto, T. Initial triggering of M-phase in starfish oocytes: A possible novel component of maturation-promoting factor besides cdc2 kinase. 125.
Olson, E.N. *See* Li, Miano, Mercer, and Olson.
Olwin, B.B. *See* Hannon, Kudla, McAvoy, Clase, and Olwin.
Oorschot, V. *See* Stoorvogel, Oorschot, and Geuze.
Orlowski, J. *See* Fukushima, Waddell, Grinstein, Goss, Orlowski, and Downey.
Ostap, E.M., and Pollard, T.D. Biochemical kinetic characterization of the *Acanthamoeba* myosin-I ATPase. 1053.

P

Pahl, H.L., Sester, M., Burgert, H.-G., and Baeuerle, P.A. Activation of transcription factor NF-κB by the adenovirus E3/19K protein requires its ER retention. 511.
Paladini, R.D., Takahashi, K., Bravo, N.S., and Coulombe, P.A. Onset of re-epithelialization after skin injury correlates with a reorganization of keratin filaments in wound edge keratinocytes: Defining a potential role for keratin 16. 381.
Pancook, J.D. *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.
Pandiella, A. *See* Cabrera, Díaz-Rodríguez, Becker, Martín-Zanca, and Pandiella.
Parkos, C.A., Colgan, S.P., Liang, T., Nusrat, A., Bacarra, A.E., Carnes, D.K., and Madara, J.L. CD47 mediates post-adhesive events required for neutrophil migration across polarized intestinal epithelia. 437.
Paschal, B.M. *See* Echeverri, Paschal, Vaughan, and Vallee.
Pearse, A. *See* Futter, Pearse, Hewlett, and Hopkins.
Pelchen-Matthews, A. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Peloquin, J.G. *See* Wang, Peloquin, Zhai, Bulinski, and Borisy.
Petersen, B.E. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.

Pfaff, M. *See* Wennerberg, Lohikangas, Gullberg, Pfaff, Johansson, and Fässler.
Pollard, T.D. *See* Ostap and Pollard.
Porter, R.M., Leitgeb, S., Melton, D.W., Swensson, O., Eady, R.A.J., and Magin, T.M. Gene targeting at the mouse cytokeratin 10 locus: Severe skin fragility and changes of cytokeratin expression in the epidermis. 925.
Posthuma, G. *See* Rijnboutt, Jansen, Posthuma, Hynes, Schornagel, and Strous.
Prekeris, R., Mayhew, M.W., Cooper, J.B., and Terrian, D.M. Identification and localization of an actin-binding motif that is unique to the epsilon isoform of protein kinase C and participates in the regulation of synaptic function. 77.
Pringle, J.R. *See* Fares, Goetsch, and Pringle.

R

Rafuse, V.F., and Landmesser, L. Contractile activity regulates isoform expression and polysialylation of NCAM in cultured myotubes: Involvement of Ca^{2+} and protein kinase C. 969.
Rajasekaran, A.K., Hojo, M., Huima, T., and Rodriguez-Boulan, E. Catenins and zonulus occulens-1 form a complex during early stages in the assembly of tight junctions. 451.
Rapraeger, A.C. *See* Lebakken and Rapraeger.
Rashid, D.J. *See* Wedaman, Meyer, Rashid, Cole, and Scholey.
Redick, J. *See* Creutz, Snyder, Daigle, and Redick.
Reed, S.I. *See* Mondésert and Reed.
Reichardt, L. *See* Wada, Kumar, Liu, Ruoslahti, Reichardt, Marvaldi, and Kanwar.
Reichmann, E. *See* Fialka, Schwarz, Reichmann, Oft, Busslinger, and Beug.
Reid, M.S. *See* Weskamp, Krätzschmar, Reid, and Blobel.
Reisfeld, R.A. *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.
Remington, S. *See* Goulielmos, Gounari, Remington, Müller, Häner, Aebi, and Georgatos.
Rieder, C.L. *See* Khodjakov, Cole, Bajer, and Rieder.
Rijnboutt, S., Jansen, G., Posthuma, G., Hynes, J.B., Schornagel, J.H., and Strous, G.J. Endocytosis of GPI-linked membrane folate receptor-α. 35.
Riley, T. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Rodriguez-Boulan, E. *See* Rajasekaran, Hojo, Huima, and Rodriguez-Boulan.
Rohrer, J., Schweizer, A., Russell, D., and Kornfeld, S. The targeting of Lampl to lysosomes is dependent on the spacing of its cytoplasmic tail tyrosine sorting motif relative to the membrane. 565.
— *See* Schweizer, Kornfeld, and Rohrer.
Romano, J. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Rosado, M. *See* Vincent, Rosado, and Davies.
Ross, A.H., Daou, M.-C., McKinnon, C.A., Condon, P.J., Lachyankar, M.B., Stephens, R.M., Kaplan, D.R., and Wolf, D.E. The neurotrophin receptor, gp75, forms a complex with the receptor tyrosine kinase TrkA. 945.
Rossetto, O., Gorza, L., Schiavo, G., Schiavo, N., Scheller, R.H., and Montecucco, C. Vamp/synaptobrevin isoforms 1 and 2 are widely and differentially expressed in nonneuronal tissues. 167.
Rousselle, P. *See* Champliaud, Lunstrum, Rousselle, Nishiyama, Keene, and Burgeson.
Rubin, K. *See* Sundberg and Rubin.
Ruoslahti, E. *See* Wada, Kumar, Liu, Ruoslahti, Reichardt, Marvaldi, and Kanwar.
Russell, D. *See* Rohrer, Schweizer, Russell, and Kornfeld.

S

Sanan, D.A. *See* Zimmerli, Majeed, Gustavsson, Stendahl, Sanan, and Ernst.
Santini, F., and Keen, J.H. Endocytosis of activated receptors and clathrin-coated pit formation: Deciphering the chicken or egg relationship. 1025.
Sapperstein, S.K., Lupashin, V.V., Schmitt, H.D., and Waters, M.G. Assembly of the ER to Golgi SNARE complex requires Uso1p. 755.
Sato-Harada, R. *See* Hirokawa, Funakoshi, Sato-Harada, and Kanai.
Sauter, M.M., Pelchen-Matthews, A., Bron, R., Marsh, M., LaBranche, C.C., Vance, P.J., Romano, J., Haggarty, B.S., Hart, T.K., Lee, W.M.F., and Hoxie, J.A. An internalization signal in the simian immunodeficiency virus transmembrane protein cytoplasmic domain modulates expression of envelope glycoproteins on the cell surface. 795.
Schekman, R.W. *See* Lupashin, Hamamoto, and Schekman.
Scheller, R.H. *See* Rossetto, Gorza, Schiavo, Schiavo, Scheller, and Montecucco.
Schiavo, G. *See* Rossetto, Gorza, Schiavo, Schiavo, Scheller, and Montecucco.
Schiavo, N. *See* Rossetto, Gorza, Schiavo, Schiavo, Scheller, and Montecucco.
Schmitt, H.D. *See* Sapperstein, Lupashin, Schmitt, and Waters.
Scholey, J.M. *See* Wedaman, Meyer, Rashid, Cole, and Scholey.
Scholze, A. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Schornagel, J.H. *See* Rijnboutt, Jansen, Posthuma, Hynes, Schornagel, and Strous.
Schütte, K. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Schwarz, H. *See* Fialka, Schwarz, Reichmann, Oft, Busslinger, and Beug.
Schweizer, A., Kornfeld, S., and Rohrer, J. Cysteine^{34} of the cytoplasmic tail of the cation-dependent mannose 6-phosphate receptor is reversibly palmitoylated and required for normal trafficking and lysosomal enzyme sorting. 577.
— *See* Rohrer, Schweizer, Russell, and Kornfeld.
Scott, S.V., and Theg, S.M. A new chloroplast protein import intermediate reveals distinct translocation machineries in the two envelope membranes: Energetics and mechanistic implications. 63.
— *See* Oda, Scott, Hefner-Gravink, Caffarelli, and Klionsky.
Sekiai, T. *See* Okumura, Sekiai, Hisanaga, Tachibana, and Kishimoto.
Sekimata, M., Tsujimura, K., Tanaka, J., Takeuchi, Y., Inagaki, N., and Inagaki, M. Detection of protein kinase activity specifically activated at metaphase-anaphase transition. 635.
Sester, M. *See* Pahl, Sester, Burgert, and Baeuerle.
Shimizu, J. *See* Okazawa, Shimizu, Kamei, Imafuku, Hamada, and Kanazawa.
Shpetner, H., Joly, M., Hartley, D., and Corvera, S. Potential sites of PI-3 kinase function in the endocytic pathway revealed by the PI-3 kinase inhibitor, wortmannin. 595.
Siu, C.-H. *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.
Slot, J.W. *See* Glickman, Morton, Slot, Kornfeld, and Geuze.
Smit, J.J.M. *See* Dunia, Smit, van der Valk, Bloemendal, Borst, and Benedetti.
Smith, E.F., and Lefebvre, P.A. *PF16* encodes a protein with armadillo repeats and localizes to a single microtubule of the central apparatus in *Chlamydomonas* flagella. 359.
Snyder, S.L. *See* Creutz, Snyder, Daigle, and Redick.
Spiess, E. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Spring, F. *See* Bartolazzi, Nocks, Aruffo, Spring, and Stamenkovic.
Springer, T.A. *See* Dustin, Ferguson, Chan, Springer, and Golan.
Stamenkovic, I. *See* Bartolazzi, Nocks, Aruffo, Spring, and Stamenkovic.
Stendahl, O. *See* Zimmerli, Majeed, Gustavsson, Stendahl, Sanan, and Ernst.
Stephens, R.M. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.
Stierhof, Y.-D. *See* Häusler, Stierhof, Wirtz, and Clayton.
Stoorvogel, W., Oorschot, V., and Geuze, H.J. A novel class of clathrin-coated vesicles budding from endosomes. 21.
Strom, S.C. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Strous, G.J. *See* Rijnboutt, Jansen, Posthuma, Hynes, Schornagel, and Strous.

Sundberg, C., and Rubin, K. Stimulation of β_1 integrins on fibroblasts induces PDGF independent tyrosine phosphorylation of PDGF β-receptors. 741.
Sundberg, H.A. *See* Friedman, Sundberg, Huang, and Davis.
Sutoh, K. *See* Aizawa, Sutoh, and Yahara.
Swanson, J.A. *See* Oh and Swanson.
Swartzman, E.E., Viswanathan, M.N., and Thorner, J. The *PALI* gene product is a peroxisomal ATP-binding cassette transporter in the yeast *Saccharomyces cerevisiae*. 549.
Swensson, O. *See* Porter, Leitgeb, Melton, Swensson, Eady, and Magin.
Sympson, C.J., Talhouk, R.S., Alexander, C.M., Chin, J.R., Clift, S.M., Bissell, M.J., and Werb, Z. Targeted expression of stromelysin-1 in mammary gland provides evidence for a role of proteinases in branching morphogenesis and the requirement for an intact basement membrane for tissue-specific gene expression: Correction. 753.

T

Tachibana, K. *See* Okumura, Sekiai, Hisanaga, Tachibana, and Kishimoto.
Takahashi, K. *See* Paladini, Takahashi, Bravo, and Coulombe.
Takeuchi, Y. *See* Sekimata, Tsujimura, Tanaka, Takeuchi, Inagaki, and Inagaki.
Taki, T. *See* Yoshida, Fujisawa-Sehara, Taki, Arai, and Nabeshima.
Talhouk, R.S. *See* Sympson, Talhouk, Alexander, Chin, Clift, Bissell, and Werb.
Tanaka, J. *See* Sekimata, Tsujimura, Tanaka, Takeuchi, Inagaki, and Inagaki.
Tang, X. *See* Benian, Tinley, Tang, and Borodovsky.
Tanimura, A., and Turner, R.J. Calcium release in HSY cells conforms to a steady-state mechanism involving regulation of the inositol 1,4,5-trisphosphate receptor Ca^{2+} channel by luminal $[Ca^{2+}]$. 607.
Tarone, G. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.
Terrian, D.M. *See* Prekeris, Mayhew, Cooper, and Terrian.
Theg, S.M. *See* Scott and Theg.
Thorner, J. *See* Swartzman, Viswanathan, and Thorner.
Tian, G. *See* Lewis, Tian, Vainberg, and Cowan.
Tinley, T.L. *See* Benian, Tinley, Tang, and Borodovsky.
Tomatis, D. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.
Tooze, S.A. *See* Dittié, Hajibagheri, and Tooze.
Tsujimura, K. *See* Sekimata, Tsujimura, Tanaka, Takeuchi, Inagaki, and Inagaki.
Turner, R.J. *See* Tanimura and Turner.

V

Vainberg, I.E. *See* Lewis, Tian, Vainberg, and Cowan.
Vallee, R.B. *See* Echeverri, Paschal, Vaughan, and Vallee.
Vance, P.J. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
van den Berghe, H. *See* Mertens, van der Schueren, van den Berghe, and David.
van der Bijl, P., Lopes-Cardozo, M., and van Meer, G. Sorting of newly synthesized galactosphingolipids to the two surface domains of epithelial cells. 813.
van der Schueren, B. *See* Mertens, van der Schueren, van den Berghe, and David.
van der Valk, M.A. *See* Dunia, Smit, van der Valk, Bloemendal, Borst, and Benedetti.
van Erck, M.G.M. *See* Koolwijk, van Erck, de Vree, Vermeer, Weich, Hanemaaijer, and van Hinsbergh.
van Hinsbergh, V.W.M. *See* Koolwijk, van Erck, de Vree, Vermeer, Weich, Hanemaaijer, and van Hinsbergh.
van Meer, G. *See* van der Bijl, Lopes-Cardozo, and van Meer.
Vaughan, K.T. *See* Echeverri, Paschal, Vaughan, and Vallee.
Vermeer, M.A. *See* Koolwijk, van Erck, de Vree, Vermeer, Weich, Hanemaaijer, and van Hinsbergh.
Vincent, I., Rosado, M., and Davies, P. Mitotic mechanisms in Alzheimer's disease? 413.
Viswanathan, M.N. *See* Swartzman, Viswanathan, and Thorner.

W

Wada, J., Kumar, A., Liu, Z., Ruoslahti, E., Reichardt, L., Marvaldi, J., and Kanwar, Y.S. Cloning of mouse integrin α_v cDNA and role of the α_v-related matrix receptors in metanephric development. 1161.
Waddell, T.K. *See* Fukushima, Waddell, Grinstein, Goss, Orlowski, and Downey.
Waddle, J.A., Karpova, T.S., Waterston, R.H., and Cooper, J.A. Movement of cortical actin patches in yeast. 861.
Walsh, K. *See* Andrés and Walsh.
Walter, P. *See* Ng and Walter.
Wang, X.M., Peloquin, J.G., Zhai, Y., Bulinski, J.C., and Borisy, G.G. Removal of MAP4 from microtubules in vivo produces no observable phenotype at the cellular level. 345.
Waters, M.G. *See* Harris and Waters.
— *See* Sapperstein, Lupashin, Schmitt, and Waters.
Waterston, R.H. *See* Waddle, Karpova, Waterston, and Cooper.
Wedaman, K.P., Meyer, D.W., Rashid, D.J., Cole, D.G., and Scholey, J.M. Sequence and submolecular localization of the 115-kD accessory subunit of the heterotrimeric kinesin-II ($KRP_{85/95}$) complex. 371.
Wegener, A.-M.K. *See* Dietrich, Neisig, Hou, Wegener, Gajhede, and Geisler.
Weich, H.A. *See* Koolwijk, van Erck, de Vree, Vermeer, Weich, Hanemaaijer, and van Hinsbergh.
Weiss, E., and Winey, M. The *Saccharomyces cerevisiae* spindle pole body duplication gene *MPS1* is part of a mitotic checkpoint. 111.
Wennerberg, K., Lohikangas, L., Gullberg, D., Pfaff, M., Johansson, S., and Fässler, R. $\beta 1$ integrin–dependent and –independent polymerization of fibronectin. 227.
Werb, Z. *See* Sympson, Talhouk, Alexander, Chin, Clift, Bissell, and Werb.
Weskamp, G., Krätzschmar, J., Reid, M.S., and Blobel, C.P. MDC9, a widely expressed cellular disintegrin containing cytoplasmic SH3 ligand domains. 717.
Whittingham, D.G. *See* Kono, Jones, Bos-Mikich, Whittingham, and Carroll.
Wickner, W. *See* Xu and Wickner.
Wigger, F. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Willard, H.F. *See* Clemson, McNeil, Willard, and Lawrence.
Winey, M. *See* Weiss and Winey.
Wirtz, E. *See* Häusler, Stierhof, Wirtz, and Clayton.
Wolf, D.E. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.

X

Xia, Y., Gil, S.G., and Carter, W.G. Anchorage mediated by integrin $\alpha 6\beta 4$ to laminin 5 (epiligrin) regulates tyrosine phosphorylation of a membrane-associated 80-kD protein. 727.
Xu, J., and Clark, R.A.F. Extracellular matrix alters PDGF regulation of fibroblast integrins. 239.
Xu, Z., and Wickner, W. Thioredoxin is required for vacuole inheritance in *Saccharomyces cerevisiae*. 787.

Y

Yagi, T. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.

Yahara, I. *See* Aizawa, Sutoh, and Yahara.
Yamamoto, T. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Yamanashi, Y. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Yasuda, M. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Yasunaga, M., Yagi, T., Hanzawa, N., Yasuda, M., Yamanashi, Y., Yamamoto, T., Aizawa, S., Miyauchi, Y., and Nishikawa, S.-I. Involvement of fyn tyrosine kinase in progression of cytokinesis of B lymphocyte progenitor. 91.
Yoshida, S., Fujisawa-Sehara, A., Taki, T., Arai, K., and Nabeshima, Y. Lysophosphatidic acid and bFGF control different modes in proliferating myoblasts. 181.

Z

Zhai, Y. *See* Wang, Peloquin, Zhai, Bulinski, and Borisy.
Zhang, J.W., and Lazarow, P.B. Peblp (Pas7p) is an intraperoxisomal receptor for the NH_2-terminal, type 2, peroxisomal targeting sequence of thiolase: Peblp itself is targeted to peroxisomes by an NH_2-terminal peptide. 325.
Zhao, X. *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.
Zhidkova, N.I. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.
Zimmerli, S., Majeed, M., Gustavsson, M., Stendahl, O., Sanan, D.A., and Ernst, J.D. Phagosome–lysosome fusion is a calcium-independent event in macrophages. 49.

SUBJECT INDEX VOLUME 132

Acanthamoeba, biochemical kinetic characterization of myosin-I ATPase of, 1053
Acetylcholine receptor, clustering of induced by agrin, mammalian muscle, requires tyrosine phosphorylation, 937
Actin
  chaperonin-mediated folding of tubulin and, 1
  movement of cortical patches of, yeast, 861
  overexpression of cofilin stimulates bundling of filaments of, membrane ruffling, cell movement, *Dictyostelium*, 335
Actin-binding motif is unique to epsilon isoform of protein kinase C, participates in regulation of synaptic function, 77
Actin-depolymerizing factor, along with cofilin is a phosphorylation-regulated protein essential for development, *Xenopus laevis*, 871
Activated receptors, endocytosis of, clathrin-coated pit formation, 1025
γ-Adaptin, AP-1 adaptor complex binds to immature secretory granules from PC12 cells, regulated by ADP-ribosylation factor, 523
Adenovirus, activation of transcription factor NF-κB by E3/19K protein in, requires ER retention, 511
Adherens junctions, estrogen-dependent c-JunER protein causes reversible loss of mammary epithelial cell polarity involving destabilization of, 1115
Adhesion
  anchorage mediated by integrin α6β4 to laminin 5 regulates tyrosine phosphorylation of membrane-associated protein, epiligrin, 727
  catenins and Zonula Occludens-1 form complex during early stages in assembly of tight junctions, 451
  CD47 mediates post-adhesive events required for neutrophil migration across polarized intestinal epithelia, 437
  cloning of mouse integrin α_v cDNA, role of α_v-related matrix receptors in metanephric development, 1161
  glycosylation of CD44 implicated in CD44-mediated cell adhesion, hyaluronan, 1199
  stimulation of β_1 integrins on fibroblasts induces PDGF independent tyrosine phosphorylation of PDGF β-receptors, 741
  tenascin-C contains distinct adhesive, anti-adhesive, and neurite outgrowth promoting sites for neurons, 681
Adhesion molecule L1, and rat homologue NILE are ligands for integrin $\alpha_v\beta_3$, human neural cell, 475
Adhesion receptors, visualization of CD2 interaction with LFA-3 and determination of two-dimensional dissociation constant for, contact area, 465
ADP-ribosylation factor, AP-1 adaptor complex regulated by, binds to immature secretory granules from PC12 cells, 523
Agrin-induced acetylcholine receptor clustering in mammalian muscle requires tyrosine phosphorylation, 937
Alzheimer's disease, mitotic mechanisms in, 413
Aminopeptidase I, identification of cytoplasm to vacuole targeting determinant in, 999
Aminopterin, import of DHFR hybrid protein into glycosomes in vivo not inhibited by, folate analogue, 311
Amnion, contains novel laminin variant, laminin 7, like laminin 6 is covalently associated with laminin 5 to promote stable epithelial–stromal attachment, human, 1189
Anaphase
  detection of protein kinase activity specifically activated at transition with metaphase, 635
  force for poleward chromosome motion in *Haemanthus* cells acts along length of chromosome during metaphase but only at kinetochore during, 1093
Annexin, identification, localization, and functional implications of an abundant, Nematode, 1079
AP-1 adaptor complex, binds to immature secretory granules from PC12 cells, regulated by ADP-ribosylation factor, 523
Apoptosis, Bcl-2 inhibits retinoic acid–induced, neural differentiation of embryonal stem cells, 955
Armadillo repeats, *PF16* encodes a protein with, localizes to single microtubule of central apparatus in *Chlamydomonas* flagella, 359
ATP hydrolysis, assembly of ER-associated protein degradation in vitro is dependent on cytosol, calnexin, and, 291
ATPase
  biochemical kinetic characterization of myosin-I ATPase, *Acanthamoeba*, 1053
  chloroplast protein import intermediate reveals distinct translocation machineries in two envelope membranes, 63
ATP-binding cassette transporter, *PALI* gene product is, peroxisomes, *Saccharomyces cerevisiae*, 549
Autocrine mechanism, differentially expressed fibroblast growth factors regulate skeletal muscle development through paracrine and, 1151
Axis formation, binding to cadherins antagonizes signaling activity of β-Catenin during, *Xenopus*, 1105
Axons, selective stabilization of Tau in, microtubule-associated protein 2C in cell bodies and dendrites, polarized localization of cytoskeletal proteins in mature neurons, 667
B lymphoblasts, biogenesis of MHC class II compartment in human I-Cell Disease, 769
B lymphocyte progenitor, cytokinesis progression of, involvement of Fyn tyrosine kinase, 91
Basic fibroblast growth factor
  cooperative effect of TNFα, VEGF, and, formation of tubular structures of human microvascular endothelial cells in fibrin matrix, role of urokinase activity, 1177
  and lysophosphatidic acid control different modes in proliferating myoblasts, 181
Bcl-2, inhibits retinoic acid–induced apoptosis during neural differentiation of embryonal stem cells, 955
Beaded-chain filaments, filensin and phakinin form, coassemble de novo in cultured cells, 643
BED1 gene, required for polarized growth and efficient bud emergence in *Saccharomyces cerevisiae*, gene encoding galactosyltransferase homologue, 137
1,2-bis-(o-Aminophenoxy)-ethane-N,N,N′,N′-tetra acetic acid, biochemically distinct steps revealed with NEM, GTPγS, and, assembly of nuclear pore, 5
Bone diseases, expression of TGF-β2 in osteoblasts results in osteoporosis-like phenotype, 195
Budding
  BED1 required for polarized growth and efficient emergence of, *Saccharomyces cerevisiae*, gene encoding galactosyltransferase homologue, 137
  from endosomes, novel class of clathrin-coated vesicles, 21
Bullous congenital ichthyosiform erythroderma, gene targeting at mouse cytokeratin 10 locus, severe skin fragility and changes of cytokeratin expression in epidermis, 925
Cadherins, binding to, antagonizes signaling activity of β-catenin during axis formation in *Xenopus*, 1105
Caenorhabditis elegans
  gene *unc-89* encodes giant modular protein composed of Ig and signal transduction domains in, required for muscle M-line assembly, 835
  identification, localization, and functional implications of an abundant Nematode annexin, 1079
  tropinin T mutations in striated and nonstriated muscle cells of, 1061
Calcium
  catenins and Zonula Occludens-1 form complex during early stages in assembly of tight junctions, 451
  cell cycle-associated change in Ca^{2+} releasing activity leads to generation of Ca^{2+} transients in mouse embryos during first mitotic division, 915
  involved in contractile activity that regulates isoform expression and polysialylation of NCAM in cultured myotubes, protein kinase C, 969
  phagosome–lysosome fusion is independent of, macrophages, 49
  release in HSY cells conforms to steady-state mechanism involving regulation of inositol 1,4,5-trisphosphate receptor Ca^{2+} channel by luminal $[Ca^{2+}]$, 607
Calnexin, assembly of ER-associated protein degradation in vitro is dependent on cytosol, ATP, and, 291
Cataracts, human multidrug resistance 3-P-glycoprotein expression in transgenic mice induces lens membrane alterations leading to, 701
Catenins
  β-catenin, binding to cadherins antagonizes signaling activity of, axis formation in *Xenopus*, 1105
  and Zonula Occludens-1 form complex during early stages in assembly of tight junctions, 451

CD2, visualization of interaction with LFA-3, determination of two-dimensional dissociation constant for adhesion receptors in contact area, 465
CD3γ, role in T cell receptor assembly, 299
CD44, glycosylation of, implicated in CD44-mediated cell adhesion to hyaluronan, 1199
CD47, mediates post-adhesive events required for neutrophil migration across polarized intestinal epithelia, 437
cdc2 kinase, novel component of maturation-promoting factor besides, triggering of M-phase in starfish oocytes, 125
Cell cycle
  myogenin expression, cell cycle withdrawal, and phenotypic differentiation are temporally separable events that precede cell fusion upon myogenesis, 657
  spindle pole body component is phosphoprotein that is modified in a manner dependent on, *Saccharomyces cerevisiae*, 903
Cell fusion, myogenin expression, cell cycle withdrawal, and phenotypic differentiation are temporally separable events that precede, myogenesis, 657
Cell movement, overexpression of cofilin stimulates bundling of actin filaments, membrane ruffling, and, *Dictyostelium*, 335
Cell spreading, syndecan-1 mediates, transfected human lymphoblastoid cells, Raji cells, 1209
Central apparatus, *PF16* localizes to single microtubule of, *Chlamydomonas* flagella, encodes a protein with armadillo repeats, 359
Chaperones, roles in protein degradation, 255
Chaperonins, mediates folding of actin and tubulin, 1
Chlamydomonas reinhardtii, *PF16* localizes to single microtubule of central apparatus in flagella of, encodes a protein with armadillo repeats, 359
Chloroplast protein import intermediate, reveals distinct translocation machineries in two envelope membranes, 63
Chromosome alignment, molecular characterization of 50-kD subunit of dynactin reveals function for complex in spindle organization and, mitosis, 617
Chromosome transport, force for poleward chromosome motion in *Haemanthus* cells acts along length of chromosome during metaphase but only at kinetochore during anaphase, 1093
c-JunER protein, causes reversible loss of mammary epithelial cell polarity involving destabilization of adherens junctions, estrogen-dependent, 1115
Clathrin-coated pit formation, endocytosis of activated receptors and, 1025
Clathrin-coated vesicles budding from endosomes, 21
Cleavage, detection of protein kinase activity specifically activated at metaphase-anaphase transition, 635
Coated pits, endocytosis of activated receptors and clathrin-coated pit formation, 1025
Cofilin
  along with actin-depolymerizing factor is a phosphorylation-regulated protein essential for development, *Xenopus laevis*, 871
  overexpression stimulates bundling of actin filaments, membrane ruffling, and cell movement in *Dictyostelium*, 335
Cryoelectron microscopy, of D1 and cytochrome b559 polypeptides, biochemical characterization, localization, two-dimensional crystals of photosystem II, 823
Crystals, two-dimensional of photosystem II, biochemical characterization, cryoelectron microscopy, localization of D1 and cytochrome b559 polypeptides, 823
Cysteine^{34}, of cytoplasmic tail of cation-dependent mannose 6-phosphate receptor is reversibly palmitoylated and required for normal trafficking and lysosomal enzyme sorting, 577
Cytochrome b559, biochemical characterization, cryoelectron microscopy, and localization of polypeptide D1 and, two-dimensional crystals of photosystem II, 823
Cytokeratins, gene targeting at mouse cytokeratin 10 locus, severe skin fragility and changes of cytokeratin expression in epidermis, 925
Cytokinesis
  actin-depolymerizing factor/cofilin is a phosphorylation-regulated protein essential for development, *Xenopus laevis*, 871
  involvement of Fyn tyrosine kinase in progression of, B lymphocyte progenitor, 91
Cytoplasm, identification of cytoplasm to vacuole targeting determinant in aminopeptidase I, 999
Cytoplasmic domain, internalization signal in, simian immunodeficiency virus transmembrane protein, modulates expression of envelope glycoproteins on cell surface, 795
Cytoplasmic tail
  cysteine^{34} of, cation-dependent mannose 6-phosphate receptor is reversibly palmitoylated and required for normal trafficking and lysosomal enzyme sorting, 577
  targeting of LAMP1 to lysomes is dependent on spacing of cytoplasmic tail tyrosine sorting motif relative to membrane, 565
Cytoskeleton, selective stabilization of Tau in axons and microtubule-associated protein 2C in cell bodies and dendrites contributes to polarized localization of proteins of, mature neurons, 667
Cytosol, assembly of ER-associated protein degradation in vitro is dependent on calnexin, ATP, and, 291
Dendrites, selective stabilization of microtubule-associated protein 2C in cell bodies and, tau in axons, polarized localization of cytoskeletal proteins in mature neurons, 667
Dictyostelium discoideum
  myosin heavy chain kinase A in, regulates myosin localization during growth and development, 101
  overexpression of cofilin stimulates bundling of actin filaments, membrane ruffling, and cell movement in, 335
Dihydrofolate reductase fusion proteins, import of into glycosomes in vivo not inhibited by folate-analogue aminopterin, 311
Disintegrin, MDC9 is widely expressed cellular, contains cytoplasmic SH3 ligand domains, 717
Dissociation constant, determination of for adhesion receptors in contact area, visualization of CD2 interaction with LFA-3, 465
Dynactin, molecular characterization of 50-kD subunit of, reveals function for complex in chromosome alignment and spindle organization during mitosis, 617
EGF, multivesicular endosomes containing internalized EGF–EGF receptor complexes mature and fuse with lysosomes, 1011
Embryogenesis, expression of *SM22α* promoter in transgenic mice, distinct transcriptional regulatory programs in vascular and visceral smooth muscle cells, 849
Embryonic stem cells, Bcl-2 inhibits retinoic acid–induced apoptosis during neural differentiation of, 955
Embryos, cell cycle-associated change in Ca^{2+} releasing activity leads to generation of Ca^{2+} transients in, mouse, first mitotic division, 915
Endocytosis
  of activated receptors and clathrin-coated pit formation, 1025
  internalization signal in simian immunodeficiency virus transmembrane protein cytoplasmic domain modulates expression of envelope glycoproteins on cell surface, 795
  of GPI-linked membrane folate receptor-α, 35
  potential sites of PI-3 kinase function in, revealed by PI-3 kinase inhibitor, wortmannin, 595
  of VAMP facilitated by synaptic vesicle targeting signal, 537
Endoplasmic reticulum
  activation of transcription factor NF-κB by adenovirus E3/19K protein requires retention in, 511
  assembly of ER to Golgi SNARE complex requires Usolp, 755
  ER membrane protein complex required for nuclear fusion, 499
Endoplasmic reticulum–associated protein degradation, assembly in vitro is dependent on cytosol, calnexin, and ATP, 291
Endosomes
  containing internalized EGF–EGF receptor complexes mature and fuse with lysosomes, 1011
  novel class of clathrin-coated vesicles budding from, 21
  potential sites of PI-3 kinase function in endocytic pathway revealed by PI-3 kinase inhibitor, wortmannin, 595
Endothelial cells, cooperative effect of TNFα, bFGF, and VEGF on formation of tubular structures of human microvascular, fibrin matrix, role of urokinase activity, 1177
Envelope membranes, chloroplast protein import intermediate reveals distinct translocation machineries in, 63
Epidermal growth factor, population expansion, clonal growth, and specific differentiation patterns in primary cultures of hepatocytes induced by HGF/SF, TGFα, and, chemically defined (HGM) medium, 1133
Epidermis, severe skin fragility and changes of cytokeratin expression in, gene targeting at mouse cytokeratin 10 locus, 925
Epithelial cells
  CD47 mediates post-adhesive events required for neutrophil migration across polarized intestinal, 437
  estrogen-dependent c-JunER protein causes reversible loss of polarity of, mammary, destabilization of adherens junctions, 1115

heparan sulfate expression in polarized, apical sorting of glypican (GPI-anchored proteoglycan) inversely related to heparan sulfate content, 487
human amnion contains novel laminin variant, laminin 7, like laminin 6 is covalently associated with laminin 5 to promote stable epithelial–stromal attachment, 1189
re-epithelialization after skin injury correlates with reorganization of keratin filaments in wound edge keratinocytes, potential role for keratin 16, 381
sorting of newly synthesized galactosphingolipids to two surface domains of, 813
Eukaryotes, roles of molecular chaperones in protein degradation, 255
Exocytosis
actin-binding motif is unique to epsilon isoform of protein kinase C, participates in regulation of synaptic function, 77
Vamp/synaptobrevin isoforms 1 and 2 are widely and differentially expressed in nonneuronal tissues, 167
Extracellular domain, role of CD3γ in T cell receptor assembly, 299
Extracellular matrix, alters PDGF regulation of fibroblast integrins, 239
Fc receptors for monomeric IgE, endocytosis of activated receptors and clathrin-coated pit formation, 1025
Fcγ receptors, role of tyrosine kinases and, Na^+/H^+ exchange activity during phagocytosis in human neutrophils, 1037
Fibrin, cooperative effect of TNFα, bFGF, and VEGF on formation of tubular structures of human microvascular endothelial cells in matrix of, role of urokinase activity, 1177
Fibroblast growth factors, regulate skeletal muscle development through autocrine and paracrine mechanisms, differentially expressed, 1151
Fibroblasts
extracellular matrix alters PDGF regulation of integrins in, 239
stimulation of β_1 integrins on, induces PDGF independent tyrosine phosphorylation of PDGF β-receptors, 741
Fibronectin, β1 integrin–dependent and –independent polymerization of, 227
Filensin, and phakinin form beaded intermediate filaments and coassemble de novo in cultured cells, 643
Flagella, *PF*16 localizes to single microtubule of central apparatus in, *Chlamydomonas*, encodes a protein with armadillo repeats, 359
Folate-analogue aminopterin, import of DHFR hybrid protein into glycosomes in vivo not inhibited by, 311
Fyn tyrosine kinase, involved in progression of cytokinesis of B lymphocyte progenitor, 91
Galactosphingolipids, sorting of newly synthesized, two surface domains of epithelial cells, 813
Galactosyltransferase, *BED1* is gene encoding homologue of, required for polarized growth and efficient bud emergence in *Saccharomyces cerevisiae*, 137
Gene targeting, at mouse cytokeratin 10 locus, severe skin fragility and changes of cytokeratin expression in epidermis, 925
Glycoproteins, internalization signal in simian immunodeficiency virus transmembrane protein cytoplasmic domain modulates expression of envelope, cell surface, 795
Glycosylation, of CD44 implicated in CD44-mediated cell adhesion to hyaluronan, 1199
Glycosyl-phosphatidyl-inositol–linked membrane folate receptor-α, endocytosis of, 35
Glypican, apical sorting of, GPI-anchored proteoglycan, inversely related to heparan sulfate content, heparan sulfate expression in polarized epithelial cells, 487
Golgi apparatus
assembly of ER to Golgi SNARE complex requires Uso1p, 755
localization of yeast early Golgi mannosyltransferase involves retrograde transport, Och1p, 985
Golgi membranes, targeting and fusion of ER-derived transport vesicles with purified, yeast, 277
gp75, neurotrophin receptor, forms complex with receptor tyrosine kinase TrkA, 945
GTPγS, biochemically distinct steps revealed with NEM, BAPTA, and, assembly of nuclear pore, 5
Gycosomes, import of DHFR hybrid protein into, in vivo, not inhibited by folate-analogue aminopterin, 311
Haemanthus, force for poleward chromosome motion in cells of, acts along length of chromosome during metaphase but only at kinetochore during anaphase, 1093
Heat-sensitive mutant, troponin T mutations in striated and nonstriated muscle cells of *Caenorhabditis elegans*, 1061
Heparan sulfate, expression in polarized epithelial cells, apical sorting of glypican (GPI-anchored proteoglycan) inversely related to heparan sulfate content, 487
Hepatocyte growth factor/scatter factor, population expansion, clonal growth, and specific differentiation patterns in primary cultures of hepatocytes induced by EGF, TGFα, and, chemically defined (HGM) medium, 1133
Hepatocytes, population expansion, clonal growth, and specific differentiation patterns in primary cultures of, induced by HGF/SF, EGF, and TGFα in chemically defined (HGM) medium, 1133
HSY cells, calcium release in, conforms to steady-state mechanism involving regulation of inositol 1,4,5-trisphosphate receptor Ca^{2+} channel by luminal $[Ca^{2+}]$, 607
Human, multidrug resistance 3–P-glycoprotein of, expression in transgenic mice induces lens membrane alterations leading to cataract, 701
Hyaluronan, glycosylation of CD44 implicated in CD44-mediated cell adhesion to, 1199
I-Cell Disease, biogenesis of MHC class II compartment in B lymphoblasts of, 769
Immature secretory granules, AP-1 adaptor complex binds to, PC12 cells, regulated by ADP-ribosylation factor, 523
Immunoglobulin domains, gene *unc-89* encodes giant modular protein composed of domains of signal transduction and, *Caenorhabditus elegans*, required for muscle M-line assembly, 835
Inflammatory diseases, CD47 mediates post-adhesive events required for neutrophil migration across polarized intestinal epithelia, 437
Inositol 1,4,5-trisphosphate, calcium release in HSY cells conforms to steady-state mechanism involving regulation of, receptor Ca^{2+} channel, luminal $[Ca^{2+}]$, 607
Integrins
$\alpha 6\beta 4$, anchorage mediated by, to laminin 5 (epiligrin), regulates tyrosine phosphorylation of membrane-associated protein, 727
$\alpha_v\beta_3$, human neural cell adhesion molecule L1 and rat homologue NILE are ligands for, 475
β_1, stimulation of, fibroblasts, induces PDGF independent tyrosine phosphorylation of PDGF β-receptors, 741
β1 integrin–dependent and –independent polymerization of fibronectin, 227
β1D, displaces β1A isoform in striated muscles, localization at junctional structures and signaling potential in nonmuscle cells, 211
cloning of mouse integrin α_v cDNA, role of α_v-related matrix receptors in metanephric development, 1161
extracellular matrix alters PDGF regulation of, fibroblast, 239
Intermediate filaments, filensin and phakinin form beaded, coassemble de novo in cultured cells, 643
Interphase, XIST RNA paints the inactive X chromosome in, novel RNA involved in nuclear/chromosome structure, 259
Karyogamy, ER membrane protein complex required for nuclear fusion, 499
Keratin
nontetrameric species is major soluble form of, *Xenopus* oocytes and rabbit reticulocyte lysates, 153
re-epithelialization after skin injury correlates with reorganization of filaments of, wound edge keratinocytes, potential role for keratin 16, 381
Kinesin II, sequence and localization of 115-kD accessory subunit of heterotrimeric complex of, $KRP_{85/95}$, 371
Kinetochore, force for poleward chromosome motion in *Haemanthus* cells acts along length of chromosome during metaphase but during anaphase only at, 1093
$KRP_{85/95}$, sequence and localization of 115-kD accessory subunit of heterotrimeric kinesin-II complex, 371
Laminin 5, anchorage mediated by integrin $\alpha 6\beta 4$ to, epiligrin, regulates tyrosine phosphorylation of membrane-associated protein, 727
Laminins, human amnion contains novel laminin variant, laminin 7, like laminin 6 is covalently associated with laminin 5 to promote stable epithelial–stromal attachment, 1189
Lamp1, targeting to lysomes, dependent on spacing of cytoplasmic tail tyrosine sorting motif relative to membrane, 565
Lens
filensin and phakinin form beaded intermediate filaments and coassemble de novo in cultured cells, 643
human multidrug resistance 3–P-glycoprotein expression in transgenic mice induces membrane alterations of, leading to cataract, 701
Ligation, human neural cell adhesion molecule L1 and rat homologue NILE are ligands for integrin $\alpha_v\beta_3$, 475
Lymphoblastoid cells, syndecan-1 mediates cell spreading in transfected human, Raji cells, 1209
Lymphocyte function-associated antigen 3, visualization of CD2 interaction with, determination of two-dimensional dissociation constant for adhesion receptors in contact area, 465

Lysophosphatidic acid, and bFGF control different modes in proliferating myoblasts, 181
Lysosomes
cysteine^{34} of cytoplasmic tail of cation-dependent mannose 6-phosphate receptor is reversibly palmitoylated and required for normal trafficking and enzyme sorting of, 577
fates of phagocytosed particles after delivery into, macrophages, 585
fusion with phagosomes, calcium-independent event in macrophages, 49
multivesicular endosomes containing internalized EGF–EGF receptor complexes mature and fuse with, 1011
targeting of LAMP1 to, dependent on spacing of cytoplasmic tail tyrosine sorting motif relative to membrane, 565
Macrophages
fates of phagocytosed particles after delivery into lysosomes of, 585
phagosome–lysosome fusion is calcium-independent event in, 49
Major histocompatibility complex class II molecules, biogenesis of, in human I-Cell Disease B lymphoblasts, 769
Mammary carcinogenesis, estrogen-dependent c-JunER protein causes reversible loss of mammary epithelial cell polarity involving destabilization of adherens junctions, 1115
Mannose 6-phosphate receptor, cysteine^{34} of cytoplasmic tail of, reversibly palmitoylated and required for normal trafficking and lysosomal enzyme sorting, 577
Mannosyltransferase, localization of, yeast early Golgi, involves retrograde transport, Och1p, 985
Matrix receptors, cloning of mouse integrin α_v cDNA, role of α_v-related matrix receptors in metanephric development, 1161
Maturation-promoting factor, novel component of, cdc2 kinase, triggering of M-phase in starfish oocytes, 125
Medium, population expansion, clonal growth, and specific differentiation patterns in primary cultures of hepatocytes in chemically defined (HGM) medium, induced by HGF/SF, EGF, and TGFα, 1133
Meiotic checkpoint, triggering of M-phase in starfish oocytes, novel component of maturation-promoting factor besides cdc2 kinase, 125
Melanoma, human neural cell adhesion molecule L1 and rat homologue NILE are ligands for integrin $\alpha_v\beta_3$, 475
Membrane folate receptor-α, endocytosis of GPI-linked, 35
Membrane proteins
activation of transcription factor NF-κB by adenovirus E3/19K protein requires ER retention, 511
of ER required for nuclear fusion, 499
Membrane ruffling, overexpression of cofilin stimulates bundling of actin filaments, cell movement, and, *Dictyostelium*, 335
Membrane trafficking, identification, localization, and functional implications of an abundant Nematode annexin, 1079
Metalloprotease/disintegrin/cysteine-rich protein 9, is widely expressed cellular disintegrin containing cytoplasmic SH3 ligand domains, 717
Metanephroi, cloning of mouse integrin α_v cDNA, role of α_v-related matrix receptors in metanephric development, 1161
Metaphase
detection of protein kinase activity specifically activated at transition with anaphase, 635
force for poleward chromosome motion in *Haemanthus* cells acts along length of chromosome during, but only at kinetochore during anaphase, 1093
Microtubule-associated protein 2C, selective stabilization in cell bodies and dendrites, Tau in axons, polarized localization of cytoskeletal proteins in mature neurons, 667
Microtubule-associated protein 4, removal from microtubules in vivo produces no observable phenotype at cellular level, 345
Microtubules
PF16 localizes to a single, central apparatus in *Chlamydomonas* flagella, encodes a protein with armadillo repeats, 359
removal of MAP4 from, in vivo, no observable phenotype at cellular level, 345
Mitosis
cell cycle-associated change in Ca^{2+} releasing activity leads to generation of Ca^{2+} transients in mouse embryos during first division of, 915
detection of protein kinase activity specifically activated at metaphase-anaphase transition, 635
molecular characterization of 50-kD subunit of dynactin reveals function for complex in chromosome alignment and spindle organization during, 617
removal of MAP4 from microtubules in vivo produces no observable phenotype at cellular level, 345
XIST RNA paints the inactive X chromosome in interphase, novel RNA involved in nuclear/chromosome structure, 259
Mitotic checkpoint, *Saccharomyces cerevisiae* spindle pole body duplication gene *MPS1* is part of, 111
Mitotic mechanisms, in Alzheimer's disease, 413
Motility, biochemical kinetic characterization of myosin-I ATPase, *Acanthamoeba*, 1053
MPS1 gene, of spindle pole body in *Saccharomyces cerevisiae*, part of mitotic checkpoint, 111
Multidrug resistance 3-P-glycoprotein, expression in transgenic mice induces lens membrane alterations leading to cataract, human, 701
Multivesicular bodies, multivesicular endosomes containing internalized EGF–EGF receptor complexes mature and fuse with lysosomes, 1011
Muscle
agrin-induced acetylcholine receptor clustering in mammalian, requires tyrosine phosphorylation, 937
gene *unc-89* required for assembly of muscle M-line, encodes giant modular protein composed of Ig and signal transduction domains, *Caenorhabditus elegans*, 835
Muscle cells
distinct transcriptional regulatory programs in vascular and visceral smooth muscle cells, expression of *SM22*α promoter in transgenic mice, 849
β1D integrin displaces β1A isoform in striated muscles, localization at junctional structures and signaling potential in nonmuscle cells, 211
tropinin T mutations in striated and nonstriated, *Caenorhabditis elegans*, 1061
Myoblasts
lysophosphatidic acid and bFGF control different modes in proliferating, 181
myogenin expression, cell cycle withdrawal, and phenotypic differentiation are temporally separable events that precede cell fusion upon myogenesis, 657
Myogenesis
lysophosphatidic acid and bFGF control different modes in proliferating myoblasts, 181
myogenin expression, cell cycle withdrawal, and phenotypic differentiation are temporally separable events that precede cell fusion upon, 657
Myosin-I, biochemical kinetic characterization of ATPase of, *Acanthamoeba*, 1053
Myosin II, *Dictyostelium* myosin heavy chain kinase A regulates localization of, growth and development, 101
Myosin heavy chain kinase A, regulates myosin localization during growth and development, *Dictyostelium*, 101
Myotubes, contractile activity regulates isoform expression and polysialylation of NCAM in, involvement of Ca^{2+} and protein kinase C, 969
Na^+/H^+ exchangers, activity of during phagocytosis in human neutrophils, role of Fcγ receptors and tyrosine kinases, 1037
Nematode, identification, localization, and functional implications of an abundant annexin in, 1079
Nephrogenesis, cloning of mouse integrin α_v cDNA, role of α_v-related matrix receptors in metanephric development, 1161
Nerve growth factor receptor complex, neurotrophin receptor forms complex with receptor tyrosine kinase TrkA, gp75, 945
Nerve growth factor-inducible large external glycoprotein, human neural cell adhesion molecule L1 and rat homologue NILE are ligands for integrin $\alpha_v\beta_3$, 475
N-ethylmaleimide, biochemically distinct steps revealed with GTPγS, BAPTA, and, assembly of nuclear pore, 5
Neural cell adhesion molecule, contractile activity regulates isoform expression and polysialylation of, cultured myotubes, involvement of Ca^{2+} and protein kinase C, 969
Neurite, tenascin-C contains distinct adhesive, anti-adhesive, and neurite outgrowth promoting sites for neurons, 681
Neurons, tenascin-C contains distinct adhesive, anti-adhesive, and neurite outgrowth promoting sites for, 681
Neurotrophin receptor, forms complex with receptor tyrosine kinase TrkA, gp75, 945
Neutrophils
CD47 mediates post-adhesive events required for migration of, polarized intestinal epithelia, 437
Na^+/H^+ exchange activity during phagocytosis in, human, role of Fcγ receptors and tyrosine kinases, 1037
Nuclear envelope breakdown, cell cycle-associated change in Ca^{2+} releasing activity leads to generation of Ca^{2+} transients in mouse embryos during first mitotic division, 915
Nuclear membrane fusion, ER membrane protein complex required for, 499

Nuclear pore complex, assembly of, steps revealed with NEM, GTPγS, and BAPTA, 5
Ochlp, yeast early Golgi mannosyltransferase, localization involves retrograde transport, 985
Oocytes
nontetrameric species is major soluble form of keratin in, *Xenopus*, rabbit reticulocyte lysates, 153
triggering of M-phase in, starfish, novel component of maturation-promoting factor besides cdc2 kinase, 125
Organelle inheritance, thioredoxin required for vacuole inheritance in *Saccharomyces cerevisiae*, 787
Osteoblasts, expression of TGF-β2 in, results in osteoporosis-like phenotype, 195
Osteoporosis, expression of TGF-β2 in osteoblasts results in osteoporosis-like phenotype, 195
PAL1 gene, product of is peroxisomal ATP-binding cassette transporter in *Saccharomyces cerevisiae*, 549
Palmitoylation, cysteine^{34} of cytoplasmic tail of cation-dependent mannose 6-phosphate receptor is reversibly palmitoylated and required for normal trafficking and lysosomal enzyme sorting, 577
Paracrine mechanism, differentially expressed fibroblast growth factors regulate skeletal muscle development through autocrine and, 1151
Pas7p, is intraperoxisomal receptor for NH_2-terminal, type 2, peroxisomal targeting sequence of thiolase, Peblp targeted to peroxisomes by NH_2-terminal peptide, 325
Patches, movement of cortical actin, yeast, 861
PC12 cells, AP-1 adaptor complex binds to immature secretory granules from, regulated by ADP-ribosylation factor, 523
Peblp, is intraperoxisomal receptor for NH_2-terminal, type 2, peroxisomal targeting sequence of thiolase, Peblp targeted to peroxisomes by NH_2-terminal peptide, Pas7p, 325
Peroxisomal targeting sequence type 2, Peblp is intraperoxisomal receptor for, NH_2-terminal, thiolase, Peblp targeted to peroxisomes by NH_2-terminal peptide, Pas7p, 325
Peroxisomes, *PAL1* gene product is peroxisomal ATP-binding cassette transporter in *Saccharomyces cerevisiae*, 549
PF16 gene, localizes to single microtubule of central apparatus in *Chlamydomonas* flagella, encodes a protein with armadillo repeats, 359
Phagocytosis
fates of phagocytosed particles after delivery into macrophage lysosomes, 585
Na^+/H^+ exchange activity during, human neutrophils, role of Fcγ receptors and tyrosine kinases, 1037
Phagosomes
fates of phagocytosed particles after delivery into macrophage lysosomes, 585
fusion with lysosomes, calcium-independent event in macrophages, 49
Phakinin, and filensin form beaded intermediate filaments and coassemble de novo in cultured cells, 643
Phospho-epitopes, mitotic mechanisms in Alzheimer's disease, 413
Phosphoprotein, modified in cell cycle-dependent manner, spindle pole body component, *Saccharomyces cerevisiae*, 903
Phosphorylation
agrin-induced acetylcholine receptor clustering in mammalian muscle requires tyrosine, 937
anchorage mediated by integrin α6β4 to laminin 5 regulates tyrosine, epiligrin, membrane-associated protein, 727
mitotic mechanisms in Alzheimer's disease, 413
Spc42p is phosphorylated component of spindle pole body with essential function during SPB duplication, *Saccharomyces cerevisiae*, 887
spindle pole body component is phosphoprotein that is modified in cell cycle-dependent manner, *Saccharomyces cerevisiae*, 903
stimulation of β_1 integrins on fibroblasts induces PDGF independent tyrosine, PDGF β-receptors, 741
TrkA receptor ectodomain cleavage generates tyrosine-phosphorylated cell-associated fragment, 427
Xenopus laevis actin-depolymerizing factor/cofilin is a protein regulated by, essential for development, 871
Photosystem II, two-dimensional crystals of, biochemical characterization, cryoelectron microscopy, localization of D1 and cytochrome b559 polypeptides, 823
PI-3 kinase, potential sites of, function in endocytic pathway revealed by PI-3 kinase inhibitor, wortmannin, 595
Platelet-derived growth factor
extracellular matrix alters regulation by, fibroblast integrins, 239
stimulation of β_1 integrins on fibroblasts induces PDGF independent tyrosine phosphorylation of PDGF β-receptors, 741
Polarity
BED1 required for efficient bud emergence and polarized growth, *Saccharomyces cerevisiae*, gene encoding galactosyltransferase homologue, 137
reversible loss in mammary epithelial cells, estrogen-dependent c-JunER protein, destabilization of adherens junctions, 1115
Polymerization, of fibronectin, β1 integrin–dependent and –independent, 227
Polypeptide D1, biochemical characterization, cryoelectron microscopy, and localization of polypeptide cytochrome b559 and, two-dimensional crystals of photosystem II, 823
Polysialylation, contractile activity regulates isoform expression and, NCAM in cultured myotubes, involvement of Ca^{2+} and protein kinase C, 969
Prokaryotes, roles of molecular chaperones in protein degradation, 255
Protein degradation
ER-associated, in vitro assembly is dependent on cytosol, calnexin, and ATP, 291
roles of molecular chaperones in, 255
Protein E3/19K, activation of transcription factor NF-κB by, adenovirus, requires ER retention, 511
Protein folding, chaperonin-mediated folding of actin and tubulin, 1
Protein import, of DHFR hybrid protein into glycosomes in vivo not inhibited by folate-analogue aminopterin, 311
Protein kinase, detection of, specifically activated at metaphase-anaphase transition, 635
Protein kinase C
actin-binding motif is unique to epsilon isoform of, participates in regulation of synaptic function, 77
involved in contractile activity that regulates isoform expression and polysialylation of NCAM in cultured myotubes, Ca^{2+}, 969
Protein transport
chloroplast protein import intermediate reveals distinct translocation machineries in two envelope membranes, 63
targeting and fusion of ER-derived transport vesicles with purified yeast Golgi membranes, 277
Proteoglycan, GPI-anchored, heparan sulfate expression in polarized epithelial cells, apical sorting of glypican inversely related to heparan sulfate content, 487
Proteolytic cleavage, TrkA receptor ectodomain cleavage generates tyrosine-phosphorylated cell-associated fragment, 427
Rabbit, nontetrameric species is major soluble form of keratin in reticulocyte lysates of, *Xenopus* oocytes, 153
Raji cells, syndecan-1 mediates cell spreading in transfected human, lymphoblastoid cells, 1209
Reticulocyte lysates, nontetrameric species is major soluble form of keratin in, rabbit, *Xenopus* oocytes, 153
Retinoic acid, Bcl-2 inhibits apoptosis induced by, neural differentiation of embryonal stem cells, 955
Retrograde transport, localization of yeast early Golgi mannosyltransferase involves, Ochlp, 985
Saccharomyces cerevisiae
BED1 required for polarized growth and efficient bud emergence in, gene encoding galactosyltransferase homologue, 137
identification of cytoplasm to vacuole targeting determinant in aminopeptidase I, 999
identification of developmentally regulated septin and involvement of septins in spore formation in, 399
localization of yeast early Golgi mannosyltransferase involves retrograde transport, Ochlp, 985
PAL1 gene product is peroxisomal ATP-binding cassette transporter in, 549
Spc42p is phosphorylated component of spindle pole body in, essential function during SPB duplication, 887
spindle pole body component in, phosphoprotein is modified in cell cycle-dependent manner, 903
spindle pole body duplication gene MPS1 is part of mitotic checkpoint in, 111
thioredoxin required for vacuole inheritance in, 787
Sec63p, ER membrane protein complex required for nuclear fusion, 499
Septins, involvement in spore formation in *Saccharomyces cerevisiae*, identification of developmentally regulated septin, 399
Signal transduction
gene *unc-89* encodes giant modular protein composed of domains of Ig and, required for muscle M-line assembly, *Caenorhabditus elegans*, 835
β1D integrin displaces β1A isoform in striated muscles, localization at junctional structures and signaling potential in nonmuscle cells, 211

Signaling
binding to cadherins antagonizes, β-catenin, axis formation in *Xenopus*, 1105
endocytosis of VAMP facilitated by synaptic vesicle targeting signal, 537
Simian immunodeficiency virus, internalization signal in, transmembrane protein cytoplasmic domain, modulates expression of envelope glycoproteins on cell surface, 795
Skeletal muscle, differentially expressed fibroblast growth factors regulate development of, autocrine and paracrine mechanisms, 1151
Skin
re-epithelialization after injury to, correlates with reorganization of keratin filaments in wound edge keratinocytes, potential role for keratin 16, 381
severe fragility of, changes of cytokeratin expression in epidermis, gene targeting at mouse cytokeratin 10 locus, 925
SM22α, expression of promoter of in transgenic mice, distinct transcriptional regulatory programs in vascular and visceral smooth muscle cells, 849
Smooth muscle cells, distinct transcriptional regulatory programs in vascular and visceral, expression of *SM22α* promoter in transgenic mice, 849
Soluble *N*-ethylmaleimide-sensitive fusion attachment protein receptor, assembly of ER to Golgi SNARE complex requires Usolp, 755
Sorting
apical sorting of glypican (GPI-anchored proteoglycan) inversely related to heparan sulfate content, heparan sulfate expression in polarized epithelial cells, 487
of newly synthesized galactosphingolipids to two surface domains of epithelial cells, 813
targeting of LAMP1 to lysomes is dependent on spacing of cytoplasmic tail tyrosine sorting motif relative to membrane, 565
Spc42p, phosphorylated component of spindle pole body with essential function during SPB duplication, *Saccharomyces cerevisiae*, 887
Spindle pole body
110-kD component of, phosphoprotein is modified in cell cycle-dependent manner, *Saccharomyces cerevisiae*, 903
duplication gene MPS1 of, part of mitotic checkpoint, *Saccharomyces cerevisiae*, 111
Spc42p is phosphorylated component of, essential function during SPB duplication, *Saccharomyces cerevisiae* 887,
Spindles, molecular characterization of 50-kD subunit of dynactin reveals function for complex in chromosome alignment and organization of, mitosis, 617
Spore formation, identification of developmentally regulated septin and involvement of septins in, *Saccharomyces cerevisiae*, 399
Src homology 3 ligand domains, MDC9 is widely expressed cellular disintegrin containing, cytoplasmic, 717
Starfish, triggering of M-phase in oocytes of, novel component of maturation-promoting factor besides cdc2 kinase, 125
Stroma, human amnion contains novel laminin variant, laminin 7, like laminin 6 is covalently associated with laminin 5 to promote stable epithelial–stromal attachment, 1189
Strongylocentrotus purpuratus, sequence and localization of 115-kD accessory subunit of heterotrimeric kinesin-II complex, $KRP_{85/95}$, 371
Synaptic vesicles
actin-binding motif participates in function of, epsilon isoform of protein kinase C, 77
endocytosis of VAMP facilitated by targeting signal of, 537
Syndecan-1, mediates cell spreading in transfected human lymphoblastoid cells, Raji cells, 1209
T cell receptor, role of CD3γ in assembly of, 299
Targeting
and fusion of ER-derived transport vesicles with purified yeast Golgi membranes, 277
identification of cytoplasm to vacuole targeting determinant in aminopeptidase I, 999
Tau, selective stabilization in axons, microtubule-associated protein 2C in cell bodies and dendrites, polarized localization of cytoskeletal proteins in mature neurons, 667
Tenascin-C, contains distinct adhesive, anti-adhesive, and neurite outgrowth promoting sites for neurons, 681
Thiolase, Peblp is intraperoxisomal receptor for NH_2-terminal, type 2, peroxisomal targeting sequence of, Peblp targeted to peroxisomes by NH_2-terminal peptide, Pas7p, 325
Thioredoxin, required for vacuole inheritance in *Saccharomyces cerevisiae*, 787
Tight junctions, catenins and Zonula Occludens-1 form complex during early stages in assembly of, 451
Tissue distribution, Vamp/synaptobrevin isoforms 1 and 2 are widely and differentially expressed in nonneuronal tissues, 167
Transcription factor NF-*x*B, activated by adenovirus E3/19K protein, requires ER retention, 511
Transforming growth factor
α, population expansion, clonal growth, and specific differentiation patterns in primary cultures of hepatocytes induced by HGF/SF, EGF, and, chemically defined (HGM) medium, 1133
β2, expression of in osteoblasts results in osteoporosis-like phenotype, 195
trans-Golgi network
biogenesis of MHC class II compartment in human I-Cell Disease B lymphoblasts, 769
sorting of newly synthesized galactosphingolipids to two surface domains of epithelial cells, 813
Transmembrane domain, role of CD3γ in T cell receptor assembly, 299
Transmembrane protein, internalization signal in cytoplasmic domain of, simian immunodeficiency virus, modulates expression of envelope glycoproteins on cell surface, 795
Triton X-100, endocytosis of GPI-linked membrane folate receptor-α, 35
TrkA receptor, TrkA receptor ectodomain cleavage generates tyrosine-phosphorylated cell-associated fragment, 427
Troponin T, mutations of in striated and nonstriated muscle cells of *Caenorhabditis elegans*, 1061
Tubulin, chaperonin-mediated folding of actin and, 1
Tumor necrosis factor-α, cooperative effect of bFGF, VEGF, and, formation of tubular structures of human microvascular endothelial cells in fibrin matrix, role of urokinase activity, 1177
Tyrosine
agrin-induced acetylcholine receptor clustering in mammalian muscle requires phosphorylation of, 937
anchorage mediated by integrin α6β4 to laminin 5 regulates phosphorylation of, epiligrin, membrane-associated protein, 727
stimulation of β_1 integrins on fibroblasts induces PDGF independent phosphorylation of, PDGF β-receptors, 741
targeting of LAMP1 to lysomes is dependent on spacing of cytoplasmic tail tyrosine sorting motif relative to membrane, 565
TrkA receptor ectodomain cleavage generates tyrosine-phosphorylated cell-associated fragment, 427
Tyrosine kinase
role of Fcγ receptors and, Na^+/H^+ exchange activity during phagocytosis in human neutrophils, 1037
TrkA, neurotrophin receptor forms complex with, gp75, 945
unc-89 gene, encodes giant modular protein composed of Ig and signal transduction domains, *Caenorhabditus elegans*, required for muscle M-line assembly, 835
Urokinase, role of, cooperative effect of TNFα, bFGF, and VEGF on formation of tubular structures of human microvascular endothelial cells in fibrin matrix, 1177
Usolp, assembly of ER to Golgi SNARE complex requires, 755
Vacuoles
identification of cytoplasm to vacuole targeting determinant in aminopeptidase I, 999
thioredoxin required for inheritance of, *Saccharomyces cerevisiae*, 787
VAMP, endocytosis of, facilitated by synaptic vesicle targeting signal, 537
Vascular endothelial cell growth factor, cooperative effect of TNFα, bFGF, and, formation of tubular structures of human microvascular endothelial cells in fibrin matrix, role of urokinase activity, 1177
Vesicle-associated membrane protein/synaptobrevin, isoforms 1 and 2 of, widely and differentially expressed in nonneuronal tissues, 167
Vesicles
novel class of clathrin-coated, budding from endosomes, 21
targeting and fusion of ER-derived transport vesicles with purified yeast Golgi membranes, 277
Wortmannin, potential sites of PI-3 kinase function in endocytic pathway revealed by, PI-3 kinase inhibitor, 595
Wound healing, re-epithelialization after skin injury correlates with reorganization of keratin filaments in wound edge keratinocytes, potential role for keratin 16, 381
Wounding, extracellular matrix alters PDGF regulation of fibroblast integrins, 239
X chromosome, XIST RNA paints the inactive, interphase, novel RNA involved in nuclear/chromosome structure, 259

Xenopus
binding to cadherins antagonizes signaling activity of β-Catenin during axis formation in, 1105
nontetrameric species is major soluble form of keratin in oocytes of, rabbit reticulocyte lysates, 153
Xenopus laevis, actin-depolymerizing factor/cofilin in, a phosphorylation-regulated protein essential for development, 871
XIST gene, paints the inactive X chromosome in interphase, novel RNA involved in nuclear/chromosome structure, 259
Yeast
movement of cortical actin patches in, 861
targeting and fusion of ER-derived transport vesicles with purified Golgi membranes of, 277
Zonula Occludens-1, and catenins form complex during early stages in assembly of tight junctions, 451